

SquaresR

View the online version at

● functions.wolfram.com

Download the

● PDF File

Notations

Traditional name

Sum of squares function

Traditional notation

$r_m(n)$

Mathematica StandardForm notation

SquaresR[m, n]

Primary definition

13.12.02.0001.01

$r_m(n)$

For nonnegative integer n , the function $r_m(n)$ is the number of representations of n as a sum of m squares of different positive or negative integers.

In particular, $r_0(n) = 0$.

Example: $r_2(2) = 4$ because $2 = 1^2 + 1^2 = (-1)^2 + 1^2 = 1^2 + (-1)^2 = (-1)^2 + (-1)^2$.

Copyright

This document was downloaded from functions.wolfram.com, a comprehensive online compendium of formulas involving the special functions of mathematics. For a key to the notations used here, see <http://functions.wolfram.com/Notations/>.

Please cite this document by referring to the functions.wolfram.com page from which it was downloaded, for example:

<http://functions.wolfram.com/Constants/E/>

To refer to a particular formula, cite functions.wolfram.com followed by the citation number.

e.g.: <http://functions.wolfram.com/01.03.03.0001.01>

This document is currently in a preliminary form. If you have comments or suggestions, please email comments@functions.wolfram.com.

© 2001-2008, Wolfram Research, Inc.